

PLANT PARASITIC NEMATODES ASSOCIATED WITH SPICES

P. SUNDARARAJU, P. K. KOSHY, AND V. K. SOSAMMA

Central Plantation Crops Research Institute Regional, Station, Kayangulam,
Krishnapuram-690 533, Kerala, India.

ABSTRACT

A literature review of the nematodes associated with the important spice plants is given. As many as 35 nematodes have been recorded on black pepper, 14 on ginger, 12 on cardamom, 6 on turmeric, 10 on cloves, 9 on nutmeg, and 6 on cinnamon. Nematodes are particularly important on black pepper, cardamom, and ginger.

India has been known since time immemorial, as the "home of spices". No detailed nematological investigations have been done on any of them in India except for a few reports of root-knot nematode infestations. In this review, an attempt has been made to consolidate the scattered information on nematodes of various spices.

1. Black pepper (*Piper nigrum* L.)

Meloidogyne sp. was the first nematode to be recorded on black pepper (Delacriox, 1902 in Cochin-China). In 1906, Butler reported root-knot nematode from black pepper in Wynad, Kerala, India. The burrowing nematode, *Radopholus similis*, is notorious for having wiped out 22 million trees (a pepper 'tree' is a teakwood or concrete post set in the field with two pepper vines entwined about it) during 20 years' time in the Bangka Island of Indonesia (Christie, 1959). Association of *R. similis* with yellows disease of pepper was first reported by Van der Vecht (1950), who made extensive field studies and also demonstrated the pathogenicity of this nematode under laboratory conditions. Later, Hubert (1957) reported the endemic occurrence of burrowing nematode in various areas of Bangka

Island and suggested a fungus-nematode complex as the cause of yellows disease. The first indication of yellows disease is the appearance of occasional yellowed leaves, which increase in numbers until, within a large portion, or even all of the foliage, may become affected. Growth ceases soon after yellowing becomes apparent, and production of pepper rapidly declines. Severe die-back and eventual death of the plants follow. The rootlets of affected vines show small lesions. These enlarge and disintegrate the rootlets. Thus, the main roots become devoid of small feeder roots and the larger laterals gradually develop necrosis also. Christie (1957) reported that pepper planted in isolated areas escaped the burrowing nematode infestation for at least five years. According to him, application of different organic and commercial fertilisers failed to retard development of yellows disease. No resistance was located in any of the pepper cultivars available then in Bangka. Recently Ichinohe (1976b) has reported an increase in area under black pepper in Bangka island during the past 20 years in newly cleared areas. According to him, the "yellows disease" attributed to *R. similis* was rare, but *Meloidogyne* sp. was widely distributed.

He differentiated the symptom picture in pepper due to *Meloidogyne* sp. and *R. similis*. Plants infested with *Meloidogyne* sp. exhibited dense yellowish discolouration of leaves, whereas those infested with *R. similis* exhibited pale yellow or whitish discoloration and the hanging yellow leaves curl inwards. Holliday and Mowat (1963) observed that infestation by *Meloidogyne* sp. did not significantly enhance the susceptibility of pepper vine to foot-rot in Sarawak. Winto (1972) studied the effect of *M. incognita* and *M. javanica* on the pepper cultivar Kuching in Malaysia and reported increased susceptibility of root-knot affected plants to infection by *Phytophthora* and other pathogens. Ting (1975) reported *Meloidogyne* spp. as the most important group of nematodes in Malaysia, causing gradual decline of black pepper characterized by unthrifty growth and yellowing of leaves. Black pepper decline in Brunei has also been reported to be associated with *Meloidogyne* and certain root rotting fungi (Anonymous, 1974b). In Brazil, Ichinohe (1975) observed that as much as 91% of pepper plants in fields near Belem, Para, were infested with *M. incognita*. Vines upto five years age were less affected than older ones. *R. similis* was not found, although symptoms of yellowing leaves similar to those attributed to *R. similis* in Indonesia were seen here also.

Though as early as in 1906, Butler, and later Ayyar (1926), had reported root-knot nematode from pepper in Wynad, Kerala, India, no follow up investigations were carried out until recently to assess the role of *M. incognita* or *R. similis* in the incidence of slow and quick wilt diseases of pepper in India. Seshadri (1972, personal communication) recorded the association of *Meloidogyne* sp. and *R. similis* with slow wilt of pepper. Venkitesan (1972) recorded high populations of *R. similis* in the rhizosphere of pepper (also Koshy, Sundararaju, and

Sosamma, 1978). Samples collected from the root system of the hybrid cultivar Panniyur-I trained on arecanut palms in Bandadka, Kasaragod, Kerala yielded very high populations (94 nematodes/g of root) of *R. similis* (Koshy et al., 1978). On young feeder roots, brown lesions could be clearly seen, followed by blackening and disintegration of the root system. The optimum temperature for extraction of nematodes from infested pepper roots was determined to be 10-20°C.

Venkitesan and Setty (1978) screened 18 cultivars of black pepper, four *Piper* species and five wild *Piper* collections against *R. similis*. The wild collection VTP 430, *Piper hymenophyllum*, and *P. attenuatum* recorded least (less than 30%) root reduction and minimum (1.5 times) nematode reproduction ninety days after inoculation. The hybrid pepper cultivar Panniyur-1 recorded 91.4% root reduction and 7.6 times nematode reproduction. Koshy and Sundararaju (1979) screened seven popular cultivars of black pepper against *M. incognita* and found the hybrid cultivar Panniyur-1 as the most susceptible one and Valiakanakadan the least susceptible.

In the Panniyur-1 seedling distribution programme, methyl bromide fumigation of nursery potting mixture under polythene cover for 24-48 hr has been recommended as a guard against the incidence of root-knot nematode in Kerala (Anonymous, 1973).

Venkitesan and Setty (1977) studied pathogenicity of *R. similis* to black pepper using 10, 100, 1000, and 10,000 nematodes as initial inoculum levels. There was decline in growth of plants with increase in inoculum levels. From the above studies, they reported that *R. similis* could be a primary incitant of slow wilt disease. Koshy et al. (1979) also studied the pathogenicity of *M. incognita* on black pepper. Even 10 larvae/cutting reduced the plant growth by 16% over a period of one year, and a maximum of 50%

reduction was noted at 1,00,000 inoculum level.

Koshy, Sosamma, and Sundararaju (1977) screened 19 species of plants commonly used as pepper standards against *M. incognita* and recommended *Garuga pinnata* for large scale use. Wherever *G. pinnata* is not available, they suggested the use of *Macaranga indica* and *Erythrina indica*.

Sundararaju, Koshy, and Sosamma (1979b) surveyed the pepper gardens of Kerala and Karnataka and reported the presence of plant parasitic nematode species belonging to 17 genera from the root zone of pepper, viz., *Aphelenchoides*, *Aphelenchus*, *Criconeoides*, *Diphtherophora*, *Ditylenchus*, *Helicotylenchus*, *Hemicyclophora*, *Hoplolaimus*, *Longidorus*, *Meloidogyne incognita*, *Paratylenchus*, *Pratylenchus*, *Radopholus similis*, *Rotylenchulus reniformis*, *Rotylenchus*, *Tylenchorhynchus*, and *Xiphinema*.

Pepper is grown in Kerala usually with coconut, arecanut, banana, ginger, turmeric, etc. All these are common hosts for root-knot and burrowing nematodes. Intensive studies are therefore suggested to develop suitable cropping patterns in such gardens. Though both these nematodes have been reported on pepper, they are not always found together. The effect of either of them on the multiplication of the other as well as their combined or individual role in the incidence of diseases such as pepper decline, slow wilt, and quick wilt also needed immediate attention.

2. Cardamom (*Elettaria cardamomum* (L.) Maton)

The nematodes *M. javanica*, *M. incognita*, and *R. similis*, were recorded from cardamom in coffee growing tracts of South India by Kumar, Viswanathan, and D'Souza (1971). They also reported cardamom to be a suitable host for the multiplication of *Pratylenchus coffeae*, *M. incognita*, and *Rotylenchulus reniformis*. Other nematodes recorded on

cardamom were *Nothocriconema cardamomi*, *N. coorgi*, and *Hemicyclophora argiensis* (Khan and Nanjappa, 1972).

Koshy et al. (1976) has recorded widespread occurrence of *M. incognita* in the cardamom nurseries in the High Ranges of Kerala, India. They tested three dosages each of fensulfothion, aldicarb, and carbofuran against *M. incognita* in cardamom nurseries (Koshy et al., 1979a) and reported that two applications of fensulfothion @ 15 kg ai/ha or aldicarb @ 5 kg ai/ha, or carbofuran @ 4 kg ai/ha first at the time of transplanting and again after three months, improved plant growth and reduced nematode population considerably, but none of these treatments controlled *M. incognita* completely in cardamom nurseries.

Sundararaju et al. (1979b) have recorded *Aphelenchoides* sp., *Helicotylenchus* sp., *Hoplolaimus* sp., *M. incognita*, *Pratylenchus* sp., and *Xiphinema* sp. from the rhizosphere of cardamom in the High Ranges of Kerala, India.

3. Ginger (*Zingiber officinale* Rosc.)

Nagakura (1930) was the first worker to report a nematode, *Meloidogyne* sp. on ginger. Hart (1956) observed parasitisation by burrowing nematode. Nadakal (1964b) obtained *M. incognita* from ginger and reported that eggs and larvae failed to survive in the upper 10 cm soil during summer months of Kerala. Root knot nematode has also been reported from the rhizome and fibrous and fleshy roots from Hawaii (Huang, 1966) and Gujarat, India, (Shah and Raju, 1977). The histopathology studies revealed that the target of infection was near the xylem pole (Shah and Raju, 1977), and that the nematode induced the formation of giant cells and active division of cells surrounding the infected area. Hyperplasia of parenchyma cells was common in the infected rhizomes and roots and gall formation was conspicuous in adventitious roots. Butler and Vilsoni (1975) reported heavy infestation

of ginger by *R. similis* in Fiji and the spread of the nematode has been found to take place through infested planting materials. The nematodes enter the rhizomes and penetrate the tissues intercellularly (Vilsoni, McClure, and Butler, 1976). Heavy infestation results in destruction of tissues and formation of channels or galleries within the rhizome. Infested ginger plants exhibit stunting, chlorosis, and sparse tillering.

For controlling root knot nematodes in Fiji, hot water treatment of ginger seed material at 50°C for 10 minutes has been recommended (Anonymous, 1971). Colbran (1972) in Australia assessed the efficacy of granular nematicides such as Mocap, Nema-cur, Temik, and Vydate against *M. javanica*, and Nema-cur was found to be the most effective. It increased the rhizome yield upto 15%. Late (November) and split applications were more promising than higher rates applied early in the season. Further studies (Colbran, 1974) showed that a high level of control of root-knot nematodes was obtained with saw dust mulching upto a depth of 50-75 mm combined with an application of Nema-cur. Such a postplanting treatment with Nema-cur or oxamyl also reduced *Fusarium* yellows of ginger caused by *Fusarium oxysporum* f. sp. *zingiberi*. In Fiji, Haynes, Partridge, and Sivan (1973) recommended a control schedule for *M. javanica* involving the use of clean seed and a ginger-dalo-fallow rotation. Both *M. javanica* and *M. incognita* have been reported from ginger in Queensland (Australia) by Pegg, Moffet, and Colbran (1974) and they suggested controlling them by hot water treatment of seed materials followed by growing the crop in fumigated soil. Sukul, Das, and De (1974), while studying the action of plant extracts against *M. incognita*, found that ethanol extract of fresh ginger rhizome possessed nematicidal properties, but, since ginger extract was phytotoxic, its use was recommended only as a preplant treatment.

Agrawal, Joshi, and Haware (1974) reported increased linear growth and thickness of hyphae of *Fusarium oxysporum* f. sp. *zingiberi* when treated with extract of ginger roots infested with *M. incognita* as compared to exposure to healthy root extract.

Sundararaju, Sosamma, and Koshy (1979a) studied the pathogenic effect of *R. similis* on ginger using different levels of population, viz., 0, 10, 100, 1000, and 10,000. In general, increase in nematode population caused directly proportional reduction in growth of plants and increase in number of root lesions and rotting. The weight of root and rhizomes was also significantly reduced. For instance, rhizome weight was reduced by 74% with an initial inoculum level of 10,000 nematodes over a period of six months, and shoot weight, root weight, and rhizome weight were reduced by 43, 56, and 40% respectively, with an initial inoculum level of 10 nematodes/plant.

Sundararaju et al. (1979b) examined soil and root samples collected from the root zone of ginger in Kerala, India, and recorded the following nematode species: *Helicotylenchus* sp., *Hemicycliophora* sp., *Hoplolaimus* sp., *M. incognita*, *R. similis*, *Rotylenchulus reniformis*, and *Tylenchorhynchus* sp.

4. Turmeric (*Curcuma longa* L.)

Root-knot nematode was the first nematode to be reported on turmeric (Ayyar, 1926). Later turmeric was reported as a host for *Meloidogyne javanica* and *M. incognita* (Nirula and Kumar, 1963) and *Radopholus similis* (Koshy and Sosamma, 1975). A *Pratylenchus* sp. also was found in association with brown rot affected tissues of mature turmeric rhizomes (Sarma, Nambiar, and Nair, 1974).

Sosamma, Sundararaju, and Koshy (1979) studied the pathogenic effect of *R. similis* on turmeric using five levels of inoculum, viz., 10, 100, 1000, 10,000 and 1,00,000. *R. similis* infested rhizome exhibited small,

shallow water soaked brownish areas. Roots which showed advanced rotting and rotten roots were found to have only the epidermis. The nematode significantly decreased shoot, root, and rhizome weights, length of lamina, and number of rhizome fingers. An initial inoculum level of 10 nematodes caused 35% reduction of rhizome weight after four months and 45% reduction at the end of the season. With 10,000 nematodes, the extent of reduction in rhizome weight was 65% and 76% respectively, after four and eight months. The nematode disseminated through infested material.

Sundararaju et al. (1979b) in a preliminary survey of Kerala, India recorded *Helicotylenchus* sp., *Hoplolaimus* sp., *Meloidogyne incognita*, *Pratylenchus* sp., and *R. similis* from the soil around the root zone of turmeric.

5. Tree Spices

Meloidogyne sp. has been reported from nutmeg (Goffart, 1953; Kumar, et al., 1972; Koshy and Sosamma, 1975) and clove (Ghesquiere, 1921). Recently, Sharma and Loof (1974) examined soil and root samples collected from clove in Bahia, Brazil and

recorded the following five species: *Helicotylenchus dihystra*, *M. incognita*, *Dolichodorus* sp., *R. reniformis*, and *Trichodorus* sp., Sundararaju et al. (1979b) have recorded the following nematode species, *Helicotylenchus* sp., *Hoplolaimus* sp., *Meloidogyne* sp., *Pratylenchus* sp., *R. similis*, *R. reniformis*, *Rotylenchus* sp., *Tylenchorhynchus* sp., and *Xiphinema* sp., on nutmeg; *Caloosia* sp., *Hoplolaimus* sp., *Meloidogyne* sp., *Pratylenchus* sp., *R. reniformis*, *Helicotylenchus* sp., *Tylenchorhynchus* sp., and *Xiphinema* sp. on clove and *Criconeoides* sp., *R. reniformis*, *Helicotylenchus* sp., *Rotylenchus* sp., and *Xiphinema* sp., on cinnamon.

From the foregoing review, it is clear that nematological investigations on spice crops are still at a very elementary level only. This is so, in spite of the fact that spices are economically very important crops in certain countries of the world and nematodes cause serious losses to several of them like pepper, ginger, and cardamom. Systematic investigations are called for particularly to determine the role of nematodes in the aetiology of pepper wilts.

Table I. Nematodes associated with spices

Nematode	Locality	Reference
I. BLACK PEPPER (<i>Piper nigrum</i> L.)		
<i>Aglenchus</i> sp.	Malayasia	Reddy (1977)
<i>Alaimus</i> sp.	Brunei, Borneo	Anonymous (1975a)
<i>Aphelenchoides</i> sp.	Bahia, Brazil; Malaysia, Thailand, Karnataka, India	Sharma & Loof (1974); Reddy (1977); Sundararaju et al. (1979b)
<i>Aphelenchus avenae</i>	Brunei, Borneo	Anonymous (1975a)
<i>Aphelenchus</i> sp.	Kerala, India	Sundararaju et al. (1979b)
<i>Aporcelamellus</i> sp.	Brunei, Borneo	Anonymous (1975a)
<i>Aporcelaimus</i> sp.	-do-	-do-
<i>Criconeoides</i> sp.	Malaysia; Thailand; Kerala, India	Reddy (1977); Sundararaju et al. (1979b)
<i>Diphtherophora</i> sp.	Thailand; Kerala, India	Reddy (1977); Sundararaju et al. (1979b)
<i>Discocriconemella limitanea</i>	Bahia, Brazil	Sharma & Loof (1974)
<i>Ditylenchus</i> sp.	Sarawak, Malaysia; Kerala, India	Reddy (1977); Sundararaju et al. (1979b)
<i>Dolichodorus</i> sp.	Bahia, Brazil	Sharma & Loof (1974)
<i>Dorylaimellus</i> sp.	Brunei, Borneo	Anonymous (1975a)

<i>Helicotylenchus dihystra</i>	Bahia, Brazil	Sharma & Loof (1974)
<i>H. erythrinae</i>	Africa; Malaysia	*Luc & Quiran (1960), Reddy (1977)
<i>H. irivandranus</i>	Kerala, India	Mohandas (1975)
<i>Helicotylenchus</i> sp.	Kerala, India; Bahia, Brazil; Brunei, Borneo; Malaysia	Venkitesan (1972); Sundararaju et al. (1979b), Sharma & Loof (1974), Anonymous (1975a), Reddy (1977).
<i>Hemicriconemoides mangiferae</i>	Thailand	Reddy (1977)
<i>Hemicyclophora</i> sp.	Kerala, India,	Venkitesan (1972), Sundararaju et al. (1979b)
<i>Heterodera marioni</i>	Indonesia	Reddy (1977)
<i>H. radiculicola</i>	Kerala, India; Kampuchea; Laos, Vietnam	Ayyar (1926); Reddy (1977)
<i>Heterodera</i> (larvae)	Kerala, India	Venkitesan (1972)
<i>Hoplolaimus columbus</i>	Malaysia	Reddy (1977)
<i>H. seinhorsti</i>	Thailand	Reddy (1977)
<i>Hoplolaimus</i> sp.	Kerala, India; Brunei, Borneo	Venkitesan (1972); Sundararaju et al. (1979b); Anonymous, (1975a)
<i>Longidorus</i> sp.	Kerala, India	Sundararaju et al. (1979b)
<i>Macroposthonia onensis</i>	Brunei, Borneo	Anonymous (1975a)
<i>Meloidogyne javanica</i>	Sarawak, Malaysia	Winto (1972), Anonymous (1974b)
<i>M. incognita</i>	Kerala, India; Para, Brazil; Sarawak; Borneo	Venkitesan (1972), Koshy et al. (1977, 1979b), Koshy and Sundararaju (1979), Sundararaju et al. (1979b), Lordello & Martins (1974), Ichinohe (1975), Anonymous (1974b)
<i>Meloidogyne</i> sp.	Cochin-china; Kerala, India; Karnataka, India; Malaysia; Brunei, Borneo; Kampuchea; Thailand; Vietnam	*Delaeriox (1902); Butler (1906); D'Souza et al. (1970), Ting (1975), Anonymous (1975a), Reddy (1977)
<i>Paratylenchus leptos</i>	Nova Teutonia, Brazil	Raski (1975)
<i>Paratylenchus</i> sp.	Kerala, India	Sundararaju et al. (1979b)
<i>Pratylenchus coffeae</i>	Karnataka, India	D'Souza et al. (1970), Kumar et al. (1971)
<i>Pratylenchus</i> sp.	Karnataka, India; Malaysia	D'souza et al. (1970); Reddy (1977)
<i>Radopholus similis</i>	Bangka island, Indonesia; Karnataka, Kerala, India; Malaysia;	*Van der Vecht (1950), Hubert (1957), Christie (1957, 1959), Ichinohe (1976), D'Souza et al. (1970), Kumar et al. (1971), Venkitesan (1972), Koshy & Sosamma (1975), Koshy et al. (1978), Sundararaju et al. (1979b), Reddy (1977)
<i>Rotylenchulus reniformis</i>	Karnataka, Kerala, India; a, Bahi Brazil; Thiland	Kumar et al. (1971), Venkitesan (1972), Sundararaju et al. (1979b), Sharma & Loof (1974), Reddy (1977)
<i>Rotylenchus</i> sp.	Kerala, India	Sundararaju et al. (1979b)
<i>Scutellonema siamense</i>	Brunei, Borneo	Anonymous (1975a)
<i>Scutellonema</i> sp.	Malaysia	Reddy (1977)
<i>Trichodorus</i> sp.	Bahia, Brazil; Thailand,	Sharma & Loof (1974), Reddy (1977),
<i>Tylenchorhynchus</i> sp.	Malaysia	Reddy (1977)
<i>Xiphinema elongatum</i>	Malayaia	Reddy (1977)
<i>X. radiculicola</i>	Brunei, Borneo; Thailand	Anonymous (1975a); Reddy (1977)

<i>X. vulgare</i>	Bahia, Brazil	Sharma & Loof (1974)
<i>Xiphinema</i> sp.	Kerala, India; Malaysia	Sundararaju et al. (1979b); Reddy (1977)
II. CARDAMOM (<i>Elettaria cardamomum</i> (L.) Maton.)		
<i>Aphelenchoides</i> sp.	Kerala, India	Sundararaju et al. (1979b)
<i>Helicotylenchus</i> sp.	-do-	-do-
<i>Hemicyclophora argiensis</i>	Karnataka, India	Khan & Nanjappa (1972)
<i>Hoplolaimus</i> sp.	Kerala, India	Sundararaju, et al. (1979b)
<i>Meloidogyne incognita</i>	Karnataka, Kerala, India.	D'souza et al. (1970), Koshy et al. (1976, 1979b), Sundararaju, et al. (1979b)
<i>Meloidogyne</i> spp.	Karnataka, India	D'souza et al. (1970)
<i>Nothocriconema cardamomi</i>	-do-	Khan & Nanjappa (1972)
<i>N. coorgi</i>	-do-	-do-
<i>Pratylenchus coffeae</i>	Karnataka, India	D'souza et al. (1970); Kumar et al. (1971)
<i>Pratylenchus</i> sp.	Karnataka, Kerala; India	D'souza et al. (1970); Sundararaju et al. (1979b)
<i>Radopholus similis</i>	Karnataka, India	D'souza et al. (1970)
<i>Rotylenchulus reniformis</i>	Karnataka, India	Kumar et al. (1971)
<i>Trichodorus minor</i>	Karnataka, India	Anonymous, (1974a)
<i>Trichodorus</i> sp.	-do-	D'souza et al. (1970)
<i>Xiphinema</i> sp.	-do-	Sundararaju et al. (1979b)
III. GINGER (<i>Zingiber officinale</i> Rosc.)		
<i>Aphelenchus avenae</i>	Malaysia	Reddy (1977)
<i>Helicotylenchus dihystra</i>	Thailand	-do-
<i>H. erythrinae</i>	-do-	-do-
<i>Helicotylenchus</i> sp.	Kerala, India	Sundararaju et al. (1979b)
<i>Hemicyclophora</i> sp.	-do-	-do-
<i>Heterodera radiculicola</i>	Japan	*Nagakura (1930)
<i>Hoplolaimus</i> sp.	Kerala, India	Sundararaju et al. (1979b)
<i>Meloidogyne acrita</i>	Fiji	Reddy (1977)
<i>M. javanica</i>	USA (Plant Quarantine Interception); Queensland, Australia	*Goodey et al. (1965), *Mumford (1960); Colbran & Davis (1969); Colbran (1972), Pegg et al. (1974)
<i>M. incognita acrita</i>	USA (Plant Quarantine Interception)	*Mumford (1963)
<i>M. incognita</i>	Queensland, Australia; Hawaii; Fiji; Kerala, Madhya Pradesh, India	Colbran (1958, 1961, 1962, 1968); Pegg et al. (1974); Huang (1966), Haynes et al. (1973), Nadakal (1963), Memmen (1973), Sundararaju et al. (1979b); Agrawal et al. (1974);
<i>Meloidogyne</i> sp.	Madhya Pradesh, Gujarat, India; Australia; Malaysia,	Kulkarni & Jain (1965); Haware & Joshi (1972); Shah and Raju, (1977); Reddy (1977)
<i>Pratylenchus pratensis</i>	Sri Lanka	*Gadd (1939)
<i>Radopholus similis</i>	Florida, USA; Fiji; Kerala, India	Hart (1956), Butler. & Vilsoni (1975), Vilsoni et al. (1976); Reddy (1977); Koshy & Sosamma (1975), Sundararaju et al. (1979a, 1979b)

<i>Rotylenchulus reniformis</i>	Kerala, Delhi, India	Sundararaju et al. (1979b); Swarup et al. (1967)
<i>Tylenchorhynchus</i> sp.	Kerala, India	Sundararaju et al. (1979b)
<i>Tylenchus</i>	Malaysia	Reddy (1977)
<i>Xiphinema basiri</i>	Rajasthan, India;	Yadav & Varma (1967)
IV. TURMERIC (<i>Curcuma longa</i> L.)		
<i>Helicotylenchus</i> sp.	Kerala, India	Sundararaju, et al. (1979b)
<i>Hoplolaimus</i> sp.	-do-	-do-
<i>Meloidogyne javanica</i>	Bihar, India	Nirula & Kumar (1963)
<i>M. incognita</i>	Bihar, Kerala, India	Nirula & Kumar (1963), Nadakal & Thomas (1964), Sundararaju et al. (1979b)
<i>Pratylenchus</i> sp.	Kerala, India	Sarma et al. (1974) Sundararaju et al. (1979b)
<i>Radopholus similis</i>	Kerala, India	Koshy & Sosamma (1975), Sosamma et al. (1979), Sundararaju et al. (1979b)
V. CLOVE (<i>Eugenia caryophyllus</i> Thunb.)		
<i>Caloosia</i> sp.	Kerala, India	Sundararaju et al. (1979b)
<i>Dolichodorus</i> sp.	Bahia, Brazil	Sharma & Loof (1974)
<i>Helicotylenchus dihystra</i>	-do-	-do-
<i>Helicotylenchus</i> sp.	Kerala, India	Sundararaju et al. (1979b)
<i>Hoplolaimus</i> sp.	-do-	-do-
<i>Meloidogyne incognita</i>	Bahia, Brazil	Sharma & Loof (1974)
<i>Meloidogyne</i> sp.	Kerala, India	Ghesquiere (1921)*; Goodey et al. (1965); Sundararaju et al. (1979b)
<i>Pratylenchus</i> sp.	Kerala, India	Sundararaju (1979b)
<i>Rotylenchus reniformis</i>	Bahia, Brazil; Kerala, India	Sharma & Loof (1974); Sundararaju et al. (1979b)
<i>Trichodorus</i> sp.	Bahia, Brazil	Sharma & Loof (1974)
<i>Tylenchorhynchus</i> sp.	Kerala, India	Sundararaju et al. (1979b)
<i>Xiphinema</i> sp.	-do-	-do-
VI. NUTMEG (<i>Myristica fragrans</i> Houtt.)		
<i>Helicotylenchus</i> sp.	Kerala, India	Sundararaju et al. (1979b)
<i>Hoplolaimus</i> sp.	-do-	-do-
<i>Meloidogyne incognita</i>	Karnataka, India	Kumar et al. (1971)
<i>Meloidogyne</i> sp.	Kerala, India	*Goffart (1953), Goodey et al. (1965), Sundararaju et al. (1979b), Sundararaju et al. (1979b)
<i>Paratylenchus</i> sp.	-do-	Koshy & Sosamma (1975), Sundararaju et al. (1979b)
<i>Radopholus similis</i>	-do-	Sundararaju et al. (1979b)
<i>Rotylenchulus reinformis</i>	-do-	-do-
<i>Rotylenchus</i> sp.	-do-	-do-
<i>Xiphinema</i> sp.	-do-	-do-
VII. CINNAMON (<i>Cinnamomum zeylanicum</i> Nees.)		
<i>Criconeimoides</i> sp.	Kerala, India	Sundararaju et al. (1979b)
<i>Helicotylenchus</i> sp.	-do-	-do-
<i>Meloidogyne</i> sp.	-do-	*Goffart (1953) Goodery et al. (1965)
<i>Rotylenchulus reinformis</i>	Kerala, India	Sundararaju et al. (1979b)
<i>Rotylenchus</i> sp.	-do-	-do-
<i>Xiphinema</i> sp.	-do-	-do-

* Original not seen

REFERENCES

- ANONYMOUS, 1971. *Annual Report for 1971*. Dept. of Agric. Fiji, 34 pp.
- ANONYMOUS, 1974 a. *Annual Report for 1972-73*. Coffee Board, Bangalore: 269 pp.
- ANONYMOUS, 1974 b. *Annual Report of the Plant Pathologist*, 1974 Bander seri Bagawan, Brunei, Borneo, Plant Pathology Laboratory, Mile-3, Jalan Berakas. 21 pp.
- ANONYMOUS, 1974 c. *Annual Report for 1973*. Central Plantation Crops Res. Inst., Kasaragod, 262 pp. 1974.
- ANONYMOUS, 1975 a. *Annual Report of the Plant Pathologist*, 1975. Bander seri Bagawan, Brunei, Borneo, Plant Pathology Laboratory, Mile-3, Jalan Berakas. 22 pp.
- ANONYMOUS, 1975 b. *Annual Report for 1974*. Sarawak, Min. of Agric. and Community Develop., 252 pp.
- AGRAWAL, P. S., JOSHI L. K., AND HAWARE, M. P. 1974. Effect of root knot extract of ginger on *Fusarium oxysporum* f. *zingiberi* Trujillo causing yellows disease. *Curr. Sci.* 43: 752.
- AYYAR, P.N.K. 1926. A preliminary note on the root gall nematode *Heterodera radicicola* and its economic importance in South India. *Madras Agri. J.* 14: 113-118.
- BARAT, H. 1952. A study on the decline of the pepper plantations in Indo-China. *Arch. Rech. Agron. Cambodge, Laos et Vietnam* 13: 92.
- *BUTLER, E. J. 1906. The wilt disease of pigeon-pea and pepper. *Agri. J. India* 1: 25-36.
- BUTLER, L., AND VILSONI F., 1975. Potential hosts of burrowing nematode in Fiji. *Fiji Agri. J.* 37: 38-39.
- CHRISTIE, J. R. 1957. The yellows disease of pepper and spreading decline of citrus. *Plant Dis. Repr.* 41: 267-268.
- CHRISTIE, J. R. 1959. *Plant Nematodes: Their Bionomics and Control*. 250 pp. Agri. Exp. Sta., Univ. of Florida, Gainesville, U.S.A.,
- COLBRAN, R. C. 1958. Studies on plant and soil nematodes. 2. Queensland host records of root-knot nematodes (*Meloidogyne* species). *Qd. J. Agric. Sci.* 15: 101-136.
- COLBRAN, R. C. 1961. Root-knot of ginger. *Qd. Agri. J.* 87: 546-547.
- COLBRAN, R. C. 1962. Studies on root knot nematode control in ginger. *Qd. J. Agri. Sci.* 19: 279-281.
- COLBRAN, R. C. 1968. Nematode control lifts ginger yields. *Qd. Agri. J.* 94: 603-605.
- COLBRAN, R. C. 1972. Studies on root knot nematode control in ginger with non-volatile nematicides applied at and after planting. *Qd. J. Agri. Anim. Sci.* 29: 275-280.
- COLBRAN, R. C. 1974. Nematode control in ginger with nematicides, selection of planting material, and sawdust mulch. *Qd. J. Agri. Anim. Sci.* 31: 231-235.
- COLBRAN, R. C., AND DAVIS, J. J. 1969. Studies on hot-water treatment and soil fumigation for control of root-knot in ginger. *Qd. J. Agric. Anim. Sci.* 26: 439-445.
- *DELACRIOX, G. 1902. A malady affecting pepper (*Piper nigrum*) in Cochinchina. *Agri. Prat. des. Pays chands* 1: 672-680.
- D'SOUZA, G. I., KUMAR A. C., VISWANATHAN P.R.K., AND SHAMANNA H. V. 1970. Relative distribution and prevalence of plant parasitic nematodes in the coffee tracts of South Western India. *Indian Coffee* 34: 329-342.
- PREIRE, F., DAS C. O., AND MONTEIRO, A. R. 1978. Nematodes associated with *Piper nigrum* in the Amazon region. Resumos dos trabalhos científicos e conferencias, III Reuniao Brasileira de Nematologia, Sociedade Brasileira de Nematologia, et da Escola Superior de Agricultura, Mossoro, 1978. Mossoro, R. N., Brazil. (1978) 82 (Pt. Colecao Mossoroense vol. 62).
- FREIRE, F., DAS C. O., AND DOS SANTOS, A.V.P. 1978. Histopathology of the roots of pepper (*Piper nigrum* L.) parasitized by *Meloidogyne incognita* and *Fusarium solani* f. *piperis*. Resumos dos trabalhos científicos e conferencias, III Reuniao Brasileira de Nematologia, Sociedade Brasileira de Nematologia, et da Escola Superior de Agricultura, Mossoro, 1978. Mossoro, R.N., (1978) 79 (Pt. Colecao Mossoroense vol. 62).
- *GADD, C. H. 1939. A destructive root disease of tea caused by the nematode *Anguillulina pratensis*. *Tea Quart.* 12: 131-139.
- *GHESQUIERE, J. 1921. "Laboratoire d'Entomologie d'Eala (Equateur) Rapports de l'entomologiste". *Bull. Agri. Congo Belge* 12: 703-732.
- *GOFFART, H. 1953. Beobachtungen an pflanzen-schadlichen Nematoden I. *Nachr. Bl. Pflschutz dienst Stuttg.* 5: 150-153.
- *GOODEY, T. 1936. On *Anguillulina oryzae* (v. Breda de Haan, 1902) Goodey, 1932, a nematode parasite on the roots of rice *Oryza sativa* L. *J. Helminth.* 14: 107-112.

- GOODEY, J. B., FRANKLIN, M. T., AND HOOPER, D. J. 1959. "Supplement to the Nematode Parasites of Plants catalogued under their Hosts 1955-1958" 66 pp. CAB, Farnham Royal, England.
- GOODEY, J. B., FRANKLIN, M. T., AND HOOPER, D. J. 1965. *T. Goodey's The Nematode Parasites of Plants catalogued under their Hosts*. 214 pp. CAB, Farnham Royal, England.
- HART, W. H. 1956. Spreading decline of citrus caused by the burrowing nematode. *Bull. Calif. Dep. Agric.* 45: 263-267.
- HAWARE, M. P., AND JOSHI L. K. 1972. Ginger. *Tech. Bull.* No. 12, pp. 21. Dept. of Plant Pathology, JNKVV, Jabalpur.
- HAYNES, P. H., PARTRIDGE, I. J., AND SIVAN, P. 1973. Ginger production in Fiji. *Fiji Agri. J.* 35: 51-56.
- HOLLIDAY, P., AND MOWAT, W. P. 1963. Foot rot of *Piper nigrum* L. (*Phytophthora palmivora*) *Phytopath.* Paper No. 5. 62 pp. Commonw. Mycol. Inst., Surrey.
- HUANG, C. S. 1966. Host-parasitic relationships of the root-knot nematode in edible ginger. *Phytopathology* 56: 755-759.
- HUBERT, F. P. 1957. Diseases of some export crops in Indonesia. *Plant Dis. Repr.* 41: 56-64.
- *HUNT, J. 1959. List of Intercepted Plant Pests, 1958. 85 pp. U. S. Dept. Agric., Agric. Res. Service. Pl. Quarant. Div., Bethesda, Md., USA.
- ICHINOHE, M. 1975. Infestation of black pepper vines by the root-knot nematode, *Meloidogyne incognita*, at Tome'-Acu, Para, Brazil. *Jap. J. Nematol.* 5: 36-40.
- ICHINOHE, M. 1976a. Infestation of black pepper by the root knot nematode in Brazil. *JARQ* 10: 17-20.
- ICHINOHE, M. 1976b. Nematode problems of black pepper in Bangka island, Indonesia. *Nematol. News Letter* 22: 2.
- ICHINOHE, M., HAMADA, M., AND YOSIDA, T. 1977. Non-host plants of *Meloidogyne incognita* on black peppers in the Amazonian region. *J. Nematol.* 9: 271-272.
- KHAN, E., AND NANJAPPA, C. K. 1972. Four new species of criconematoidea (Nematoda) from India. *Indian J. Nematol.* 2: 59-68.
- KOSHY, P. K., AND SOSAMMA, V. K. 1975. Host range of *Radopholus similis* (Cobb, 1893) Thorne, 1949. *Indian J. Nematol.* 5: 255-257.
- KOSHY, P. K., NAIR, C.P.R., SOSAMMA, V. K. AND SUNDARARAJU, P. 1976. On the incidence of root knot nematode in cardamom nurseries. *Indian J. Nematol.* 6: 174-175.
- KOSHY, P. K., AND SUNDARARAJU, P. 1979. Response of seven black pepper cultivars to *Meloidogyne incognita*. *Nematol. medit.* (in press).
- KOSHY, P. K., SOSAMMA, V. K., AND, SUNDARARAJU, P. 1977. Screening of plants used as pepper standards against root knot nematode. *Indian Phytopath.* 30: 128-129.
- KOSHY, P. K., SUNDARARAJU, P., AND SOSAMMA, V. K. 1978. Occurrence and distribution of *Radopholus similis* (Cobb, 1893) Thorne, 1949 in South India. *Indian J. Nematol.* 8: 49-58.
- KOSHY, P. K., SUNDARARAJU, P., AND WILSON, K. I. 1979a. Efficacy of certain systemic nematicides against *Meloidogyne incognita* in a cardamom nursery. *Nematol. medit.* (in press).
- KOSHY, P. K., PREMACHANDRAN, D., SOSAMMA, V. K., AND PREMKUMAR, T. 1979b. Effect of *Meloidogyne incognita* population on black pepper (*Piper nigrum* L.) *Indian Phytopath.* (in press).
- KULKARNI, S. N., AND JAIN, A. C. 1965. *Meloidogyne* sp. in rhizomes of ginger (*Zingiber officinale* Rosc.). *Indian Phytopath.* 18: 391-392.
- KUMAR, A. C., VISWANATHAN, P.R.K., AND D'SOUZA, G. I. 1971. A study of plant parasitic nematodes of certain commercial crops in coffee tracts of South India. *Indian Coffee* 35: 222-224.
- KUMAR, A. C., VISWANATHAN, P.R.K., AND D'SOUZA, G. I. 1972. Record of *Radopholus similis* (Cobb, 1893). Thorne, 1949 and other parasitic nematodes of certain commercial crops in coffee tracts of South India. *Indian Coffee* 36: 1-3.
- *LORDELLO, L.G.E., AND SILVA, H. M. E. 1974. A nematode harmful to black pepper. *Rev. de Agri. Piracicaba* 49: 165-166.
- *LUC, M., AND DE GUIRAN, G. 1960. "Less nematodes associes aux plants del'ouest Africain" *Liste Preliminaire. Agron. Trop. Nogent* 15: 434-449.
- MAMMEN, K. V. 1973. Root gall nematodes as a serious pest of ginger in Kerala. *Curr. Sci.* 42: 549.
- MARTIN, G. C. 1956. The common root-knot nematode and crop rotation. *Rhod. Fmr.* 20: pp.
- MOHANDAS, C. 1975. *Helicotylenchus trivandranus* sp.n. (Nematoda: Hoplolamidae) from Kerala, India. *Indian J. Nematol.* 5: 105-107.
- *MUMFORD, B. C. 1960. List of Intercepted Plant Pests, 1959. 86 pp. U.S. Dept. Agric., Agric. Res. Service. Pl. Quarant. Div., Bethesda, Md., USA.
- *MUMFORD, B. C. 1963. List of Intercepted Plant Pests, 1962. 88 pp. U.S. Dept. Agric., Agric. Res. Service. Pl. Quarant. Div. Bethesda, Md., USA.
- NADAKAL, A. M. 1963. *Meloidogyne* spp. infecting certain plants in Kerala. *Curr. Sci.* 32: 360-361.
- NADAKAL, A. M. 1964a. Studies on plant parasitic nematodes of Kerala III. An additional list of plants attacked by root-knot nematode *Meloidogyne* sp. *J. Bombay Nat. Hist. Soc.* 61: 467-469.
- NADAKAL, A. M. 1964b. Studies on plant parasitic nematodes of Kerala. IV. Host-range, seasonal variations and summer fallow survival of zingiber root-knot nematode, *Meloidogyne incognita* (Koroid & White) Chitwood. *Indian J. Expl. Biol.* 2: 203-207.
- NADAKAL, A. M., AND THOMAS, N. 1964. Studies on plant parasitic nematodes of Kerala. *Curr. Sci.* 33: 247-248.
- *NAGAKURA, K. 1930. "Über den Bau und die Lebensgeschichte der *Heterodera radiculicola* (Greef) Muller". *Jap. J. Zool.* 3: 95-160.
- NAMBIAR, K.K.N., AND SARMA, Y. R. 1977. Wilt diseases of black pepper. *J. Plant. Crops* 5: 92-103.
- NIRULA, K. K., AND KUMAR, R. 1963. Collateral hosts of root-knot nematode. *Curr. Sci.* 32: 221-222.
- PEGG, K. G., MOFFETT, M. L., AND COLBRAN, R. C. 1974. Diseases of ginger in Queensland. *Qd. Agri. J.* 100: 611-618.
- RADHA, K., AND RAWTHER, T.S.S. 1976. Pests and diseases of gardenland crops. *Indian Fmg.* 25: 31-35.
- RASKI, D. J. 1975. Revision of the genus *Paratylenchus* Micoletzky, 1922 and descriptions of new species. *J. Nematol.* 7: 15-34.
- REDDY, D. B. 1977. Pests, Diseases and Nematodes of major Spices and Condiments in Asia and the Pacific. *Tech. Doc.* 108: 14, Plant Protection Committee for the South East Asia and Pacific Region, FAO, Bangkok.
- SARMA, Y. R., NAMBIAR, K.K.N., AND NAIR, C.P.R. 1974. Brown rot of turmeric. *J. Plant. Crops.* 2: 33-34.
- SHAH, J. J., AND RAJU, E. C. 1977. Histopathology of ginger (*Zingiber officinale*) infested by soil nematode *Meloidogyne* sp. *Phyton* (Austria) 16: 79-84.
- SHARMA, R. D., AND LOOF, P.A.A. 1974. Nematodes of cocoa region of Bahia, Brazil IV. Nematodes in the rhizosphere of pepper (*Piper nigrum* L.) and clove (*Eugenia caryophyllata* Thumb). *Rev. Theobroma* 4: 26-32.
- SHER, S. A., CHURAM, C., AND PHOLCHARDEN, S. 1969. Pepper yellows disease and nematodes in Thailand. *FAO Pl. Prot. Bull.* 17: 33.
- SOSAMMA, V. K., SUNDARARAJU, P., AND KOSHY P. K. 1979. Effect of *Radopholus similis* population on turmeric. *All India Nematology Symposium*. March 21-22, 1979. Orissa University of Agriculture and Technology, Bhubaneswar.
- SUKUL, N. C., DAS, P. K., AND DE, G. C. 1974. Nematicidal action of some edible crops. *Nematologica* 20: 187-191.
- SUNDARARAJU, P., SOSAMMA, V. K., AND KOSHY, P. K. 1979a. Pathogenicity of *Radopholus similis* on ginger. *All India Nematology Symposium*, March 21-22, 1979. Orissa University of Agriculture and Technology Bhubaneswar.
- SUNDARARAJU, P., KOSHY, P. K., AND SOSAMMA, V. K. 1979b. Survey of plant parasitic nematodes associated with spices in Kerala and Karnatalaka. *Second Ann. Symp. Plantation Crops*. June 1979, Ootacamund, Nilgiri Hills, Tamil Nadu.
- SWARUP, G., NATH, R.P., AND SETHI, C.L. 1967. The plant parasitic nematode genus *Rotylenchulus* in India. *Indian Phytopath.* 20: 118-123.
- TING, W. P. 1975. Plant pathology in Peninsular Malaysia. *Rev. Pl. Path.* 54: 267-305.
- *VECHT, J. VAN DER, 1950. Plant Parasitic Nematodes. 16-45 pp. in Karshoven, L.G.E. and Vecht, J.v.d. (ed.) *Diseases of Cultivated Plants in Indonesia*. Vol. I (in Dutch). W. van Hoeve, 's-Gravenhage.
- VENKITESAN, T. S. 1972. On the occurrence of plant parasitic nematodes associated with different crops in Cannanore District, Kerala. *Agri. Res. J. Kerala* 10: 179-180.
- VENKITESAN, T. S. 1976. Studies on the burrowing nematode, *Radopholus similis* (Cobb) Thorne, 1949 on pepper (*Piper nigrum* L.) and its role in slow wilt disease. Ph.D. Thesis. 122 pp. University of Agri. Sci., Bangalore, India.
- VENKITESAN, T. S. AND SETTY, K.G.H. 1977. Pathogenicity of *Radopholus similis* to black pepper (*Piper nigrum*). *Indian J. Nematol.* 7: 17-26.

- VENKITESAN, T. S. AND SETTY, K.G.H. 1978. Reaction of 27 black pepper cultivars and wild forms to the burrowing nematode. *Radopholus similis* Cobb, Thorne, *J. Plant. Crops* 6: 81-83.
- VENKITESAN, T. S. AND CHARLES, J. S. 1979. A note on the chemical control of nematodes infesting pepper vines in Kerala. *Second Ann. Symp. Plant. Crops*. June 1979. Ooctacamund, Nilgiri Hills, Tamil Nadu.
- VILSONI, F. 1974. New Records, *FAO: Q. Newsl.* (South East Asia and Pacific) 17: 4-5.
- VILSONI, F., Mc CLURE, M. A., AND BUTLER, L. D. 1976. Occurrence, host range and histopathology of *Radopholus similis* in ginger (*Zingiber officinale*). *Plant Dis. Repr.* 60: 417-420.
- WINTO, S. R. 1972. Effect of *Meloidogyne* species on the growth of *Piper nigrum* L. *Malaysian Agri. Res.* 1: 86-90.
- YADAV, B. S. AND VARMA, M. K. 1967. New host plants of *Xiphinema basiri* and *X. indicum*. *Nematologica* 13: 469.
- ZEVEN, A. C. 1976. Black pepper *Piper nigrum*. (Ed.) *Evolution In Simmonds NW of Crop Plants*. Longman, London, pp. 234-235.

*Original not seen