

See discussions, stats, and author profiles for this publication at:
<https://www.researchgate.net/publication/291355656>

Microclimatic and photosynthetic characteristics in arecanut and cardamom mixed cropping system v

Article · January 2000

CITATIONS

0

READS

5

5 authors, including:



[Ankegowda S J](#)

Indian Institute of Spices Research

25 PUBLICATIONS 14 CITATIONS

SEE PROFILE

Microclimatic and photosynthetic characteristics in arecanut and cardamom mixed cropping system

V S Korikanthimath, S J Ankegowda, N Yadukumar¹, Rajendra Hegde & M M Hosmani²

Cardamom Research Centre

Indian Institute of Spices Research

Appangala, Heravanad P. O., Madikeri - 571 201, Karnataka, India.

Received 29 June 1998; Revised 17 February 2000; Accepted 3 March 2000.

Abstract

The microclimatic and photosynthetic characteristics of arecanut (*Areca catechu*) and cardamom (*Elettaria cardamomum*) mixed cropping system were studied at Sirsi (Karnataka, India). Arecanut canopy received higher amount of light and also photosynthetically active radiation. Arecanut also recorded higher photosynthetic rate, transpiration rate, stomatal conductance, intercellular CO₂ concentration, carboxylation efficiency and water use efficiency at gas exchange level.

Key words: arecanut, *Areca catechu*, cardamom, *Elettaria cardamomum*, microclimate, mixed cropping, photosynthetic characteristics.

In mixed cropping systems plants compete for light, water, soil and space. Competition occurs when the immediate supply of a single necessary factor falls below the combined demands of the plants (Donald 1963). In multistoreyed cropping systems, the leaf canopy of the components of crop combinations occupy different vertical layers with the tallest component having foliage tolerant to strong light and high evaporative demand and the shorter component(s) having foliage requiring or tolerating shade and/or high humidity (Nair 1979). In mixed cropping systems, even if all other conditions are favourable for the growth of intercrops, light is a limiting factor since light penetration of a plant stand is reduced through interception and absorption by the taller canopy. In this situation, shade loving species which require low light intensities are likely to perform better (Kasturibai *et al.* 1991; Balasimha & Yadukumar 1993; Rethinam & Venugopal 1994). In arecanut (*Areca catechu* L.) plantations situated at higher elevations, especially in Uttara Kannada District in Karnataka (India), arecanut occupies the top most canopy and shade loving crops like cardamom (*Elettaria cardamomum* Maton) are grown under its shade. Arecanut has a fairly deep

root system which can tap water and nutrients from deeper layers, whereas cardamom is a shallow rooted crop, which can absorb water and nutrients from top layers. The microclimate and photosynthetic characteristics in arecanut and cardamom mixed cropping system was studied at Sirsi (Uttara Kannada District, Karnataka) and the results are reported here.

The field experiment was undertaken at Sirsi and arecanut (Sirsi local) was planted at 2.7 m x 2.7 m spacing in both mono and mixed cropping systems. Cardamom (CI-37, Malabar type) was introduced between rows of arecanut at a spacing of 2.7 m x 1.2 m. The experiment was laid out in a complete randomised block design with eight replications. The age of arecanut was 10 years and cardamom 3 years at the termination of the experiment. Physiological parameters namely, photosynthetic rate (A), transpiration rate (E), stomatal conductance (g_s), intercellular CO₂ concentration (C_i), and microclimatic parameters such as light interception (%), photosynthetically active radiation (PAR), relative humidity (%), leaf temperature and air temperature (°C) were measured by using leaf chamber analyzer (LCA-3) during March 1993-94 at 10-12 am on fully

¹ National Research Centre for Cashew, Puttur - 574 202, Karnataka, India.

² University of Agricultural Sciences, Dharwad - 580 005, Karnataka, India.

Table 1. Microclimatic variables in arecanut and cardamom mixed cropping system

Crop	Light interception (%)	PAR ($\mu\text{molm}^{-2}\text{sec}^{-1}$)	Relative humidity (%)	Air temperature ($^{\circ}\text{C}$)	Leaf temperature ($^{\circ}\text{C}$)
Arecanut	59.1	886.0	23.4	31.5	31.0
Cardamom	40.3	350.0	25.3	29.5	29.9
't' test	**	**	**	**	**

PAR = Photosynthetically active radiation

** = Significant at $P < 0.01$

opened intact leaves of arecanut and cardamom. The ratios of A/E, A/C_i and A/g_s were also computed. The data were analysed by 't' test.

Microclimate

All the microclimatic parameters were significantly different in mixed cropping system of arecanut and cardamom. Arecanut received higher amount of light (59.06%) compared to cardamom (40.25%) due to arecanut crown forming the top most canopy and cardamom constituting the lower canopy in the cropping system. Similar results were observed with PAR also i.e., arecanut

Table 2. Photosynthetic characteristics in arecanut and cardamom mixed cropping system

Crop	Pn ($\mu\text{molCO}_2\text{m}^{-2}\text{s}^{-1}$)	Tr ($\mu\text{molH}_2\text{Om}^{-2}\text{s}^{-1}$)	SC ($\text{molm}^{-2}\text{s}^{-1}$)	Int. CO ₂ (ppm)
Arecanut	4.2	4.8	0.20	286.0
Cardamom	4.1	4.0	0.24	316.0
't' test	NS	**	**	**

Pn = Photosynthetic rate; Tr = Transpiration rate; SC = Stomatal conductance; Int. CO₂ = Internal CO₂ concentration
NS = Non significant; ** = Significant at $P < 0.01$

received 886.0 $\mu\text{molm}^{-2}\text{sec}^{-1}$, whereas cardamom received 350.0 $\mu\text{molm}^{-2}\text{sec}^{-1}$. Relative humidity was low under arecanut canopy (23.45%) compared to cardamom (25.30%). Air and leaf temperatures were higher in arecanut than cardamom due to higher light interception (Table 1).

Physiological parameters

Transpiration rate, stomatal conductance and intercellular CO₂ concentration showed significant variations. Photosynthetic rate was higher in arecanut (4.2 $\mu\text{molCO}_2\text{m}^{-2}\text{sec}^{-1}$) compared to cardamom (4.1 $\mu\text{molCO}_2\text{m}^{-2}\text{sec}^{-1}$). Transpiration rate was also higher in arecanut (4.8 $\mu\text{molH}_2\text{Om}^{-2}$

Table 3. Mean values of A/E, A/g_s and A/C_i in arecanut and cardamom mixed cropping system

Crop	A/E	A/g _s	A/C _i
Arecanut	0.89	21.2	0.015
Cardamom	1.03	17.5	0.013
't' test	**	**	**

A = Assimilation rate; E = Transpiration rate; g_s = Stomatal conductance; C_i = Internal CO₂ concentration

** = Significant at $P < 0.01$

sec^{-1}) compared to cardamom (4.0 $\mu\text{mol H}_2\text{Om}^{-2}\text{sec}^{-1}$) due to higher light interception by the arecanut canopy (Table 2). Stomatal conductance was higher in cardamom (0.24 $\text{molm}^{-2}\text{sec}^{-1}$) than arecanut (0.20 $\text{molm}^{-2}\text{sec}^{-1}$). Intercellular CO₂ concentration was higher in cardamom (316.0 ppm) compared to arecanut (286.0 ppm). A/C_i which indicates carboxylation efficiency (Farquhar & Sharkey 1982) and A/E and A/g_s which indicate the intrinsic water use efficiency of crop plants were higher in arecanut (Table 3) which was due to higher light interception and intrinsic plant factors. The relation of 'A' and C_i has been reported to be linear in other crops (Farquhar & Sharkey 1982; Marco *et al.* 1988; Balasimha *et al.* 1991). Similar results were also observed in cashew (Balasimha & Yadukumar 1993).

References

- Balasimha D & Yadukumar N 1993 Effect of plant density on photosynthesis in cashew. *Indian J. Plant Physiol.* 36 : 5-7.
- Donald D M 1963 Competition among pasture plants. *Adv. Agron.* 15 : 1-118.
- Farquhar G D & Sharkey T D 1982 Stomatal conductance and photosynthesis. *Annu. Rev. Plant Physiol.* 33 : 317-345.
- Kasturibai K V, Voleti S R, Ramadasan A & Kailasam C 1991 Growth and dry matter production in

- component crops under monocropping and high density multi species cropping system. *J. Plantn. Crops* 18 (Suppl.) : 151-155.
- Marco G D, Massacii A & Gabrielli R 1988 Drought effect of photosynthesis and fluorescence in hard wheat cultivars grown in the field. *Physiol. Plant.* 74 : 385-390.
- Nair P K R 1979 Intensive multiple cropping with coconuts in India. *Adv. Agron. Crop Sci.* 6 : 1-12
- Rethinam P & Venugopal K 1994 Cropping systems in plantation crops. In: Chadha K L & Rethinam P. (Eds.) *Advances in Horticulture Vol. 9, Plantation and Spice Crops Part I* (pp. 605-620). Malhotra Publishing House, New Delhi.