

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/304104557>

Response of soil enzymes to elevated CO₂ and temperature in weeds associated with rice-wheat cropping system

Article · January 2016

DOI: 10.5958/0974-8164.2016.00006.X

CITATIONS

0

READS

31

4 authors:



[C. Sarathambal](#)

Indian Council of Agricultural Research

18 PUBLICATIONS 23 CITATIONS

[SEE PROFILE](#)



[Meenal Rathore](#)

Indian Council of Agricultural Research-Direc...

20 PUBLICATIONS 23 CITATIONS

[SEE PROFILE](#)



[Disha Jaggi](#)

Directorate of Weed Science Research

17 PUBLICATIONS 27 CITATIONS

[SEE PROFILE](#)



[Bhumesh Kumar](#)

Directorate of Weed Science Research

35 PUBLICATIONS 371 CITATIONS

[SEE PROFILE](#)



Response of soil enzymes to elevated CO₂ and temperature in weeds associated with rice-wheat cropping system

C. Sarathambal*, M. Rathore, D. Jaggi and Bhumesh Kumar
ICAR-Directorate of Weed Research, Jabalpur, Madhya Pradesh 482 004

Received: 26 November 2015; Revised: 14 January 2016

ABSTRACT

Biological properties of the soil have often been proposed as early and sensitive indicators of soil ecological stress or other environmental changes. In the present investigation, the soil samples were collected from weeds associated with rice-wheat cropping system from open-top chambers to assess the effect of elevated CO₂ and temperature on soil enzymes. In *Rabi* season, higher activity of FDA hydrolysis rate was noticed in wild oat with 26.5 (µg fluorescein/g/h) at elevated CO₂ + ambient temperature levels. Dehydrogenase activity was higher at enrichment of CO₂ for *P. minor* with 35.9 (µg TPF/g soil/24 h). Urease significantly was higher in enrichment of CO₂ + elevated temperature with wild oat 34.6 (µg NH₄/g soil/24 h) followed by wheat (31.5 µg NH₄/g soil/24 h) with elevated CO₂ + ambient temperature. In *Kharif* season, enrichment of CO₂ concentration in rhizosphere of *Echinochloa crusgalli* recorded with higher FDA hydrolysis rate (19.8 µg fluorescein/g/h), dehydrogenase activity (39.8 µg TPF/g soil/24 h) and urease activity (45.6 µg NH₄/g soil/24 h) respectively. We found the carbon dioxide enrichment significantly increased the soil enzymes like dehydrogenase, fluorescein diacetate (FDA) hydrolysis and urease activity in weeds rhizosphere than the crops.

Key words: Elevated CO₂, Elevated temperature, Open top chambers, Rice, Soil enzymes, Wheat

Indian agriculture has done remarkably well by increasing food production from 50 Mt in 1951 to 252 Mt in 2011-2012 assuring food security to the nation. However, perceived impacts of climate change could adversely affect the food output. Over the last two decades, global warming was easily dismissed as the natural fluctuation in an ever changing global climate. Although there exists natural fluctuations in climate, human induced changes are predominantly during the current global warming trend (Stager 2012). Carbon dioxide is one of the major contributors of greenhouse gases which, being a primary substrate for photosynthesis, may have a significant impact on plant metabolism that directly affect the overall plant growth and soil health. Soil enzyme activities are the indicators of soil microbial functioning, soil physicochemical conditions and nutrient cycling in soils. The use of soil enzymatic activities as possible indicators of changes in below ground processes is induced by CO₂ enrichment. Many factors can influence the activity of soil enzymes. Temperature changes can affect microbial diversity and enzymatic activity directly. A source of microbial substrates for enzymatic decomposition could be rhizodeposition and secretion of soluble root exudates. Microbes could have various positive and negative feedback responses to temperature. Whether, changes in microbial processes lead to a net

*Corresponding author: saratha6@gmail.com

positive and negative feedback for greenhouse gas emission is unclear. The reason is that microbes live in very diverse communities that interact with organisms and the environments in complex ways. It is important and there is an urgent need for more empirical knowledge that will fundamentally improve our ability to predict feedbacks of the terrestrial carbon cycle in agricultural ecosystems to anticipated climate change.

Climate change factors such as elevated atmospheric carbon dioxide (CO₂) can exert significant impacts on soil microorganisms and the ecosystem level processes they mediate. Therefore, the present investigation was carried out to study the effects of elevated CO₂ and temperature concentrations on soil enzyme activities in the weeds associated with rice-wheat ecosystem in open-top chambers (OTCs).

MATERIALS AND METHODS

Field experiment was laid out to study the changes of enzyme activity in vertisols under open top chambers (OTCs) to study the effect of elevated CO₂ and temperature on soil enzymes, at crop physiological maturity stage during *Kharif* and *Rabi* season (2013-14). The treatments included i) ambient CO₂ + ambient temperature ii) ambient CO₂ + elevated temperature iii) elevated CO₂ + ambient

temperature iv) elevated CO₂ + elevated temperature in open top chambers (OTCs) placed in ICAR-Directorate of Weed Research, Jabalpur. The elevated temperature was maintained by putting infrared lamps inside the OTC chambers and was controlled by the automatically calibrated temperature sensors. Air temperature and relative humidity was monitored continuously at every 5 min interval in each OTC chamber by temperature– humidity-calibrated sensors. During sampling, the crops were gently pulled out, and loose soil was shaken off the roots and the soil that adhered strongly to the roots was carefully brushed and sampled as rhizosphere soil. All samples were performed in triplicate, and were used to quantify the soil enzymes.

Dehydrogenase activity was assayed by the method of Casida *et al.* (1964). Moist soil samples (4 g) were placed in 16 x 150 mm² test tubes to which was added 1 ml of 3% aqueous solution of 2,3,5-triphenyl tetrazolium chloride, 40 mg CaCO₃ and 2.5 ml distilled water. The contents of each tube were then mixed with a glass rod and incubated for 24 h at 37°C. Triphenyl formazan (TPF) was extracted by transferring the soil with the aid of methanol from each tube to a funnel plugged with absorbent cotton and the colour intensity determined in a spectrophotometer at a wave length of 485 nm. The dehydrogenase activity was expressed as µg TPF formed/g soil/24h.

The determination of fluorescein diacetate (FDA) hydrolysis (Schnurer and Rosswall 1982) was carried out in 2 g of field moist soil, where 15 ml of 60 mM potassium phosphate buffer (pH 7.6) was added. 100 µl of substrate solution was added to samples. After 20 min of incubation, the reaction was stopped by adding 15 ml of chloroform: methanol (2: 1). After filtering the solutions (Whatman No. 42), absorbance was measured at 490 nm on a UV Visible Spectrophotometer and calculated from the standard graph of fluorescein. The FDA hydrolysis rate was expressed as µg fluorescein/g/h.

Urease activity was assayed by the method of Bremner and Mulvaney (1978). Ten gram of dry and sieved soil was taken in a 100 ml volumetric flask. To this 1.5 ml of toluene was added, mixed well and incubated for 15 min. Then 10 ml of 10% urea solution and 20 ml of citrate buffer were added, mixed thoroughly, stoppered and incubated for 3 h at 37° C. Then the volume was made up to 100 ml with distilled water, mixed by shaking immediately. The contents were filtered through Whatman No.1 filter paper and 1ml of filtrate was pipetted out into 50 ml volumetric flask. To this 9 ml of distilled water, 4 ml

of phenate and 3 ml of NaOCl were added, mixed well and allowed to stand for 20 min. The volume was made up to 50 ml and mixed well. The bluish green colour developed was read at 630 nm. The concentration of urease in the sample was obtained from the standard graph using diammonium sulphate. Urease activity was expressed µg NH₄/g soil/24h.

All the data were subjected to statistical analysis with softwares, SPSS (Kirkpatrick and Feenay 2005) and Microsoft Excel for Windows 2007 add-ins with XLSTAT Version 2010.5.05 (XLSTAT, 2010). Statistically significant differences between the treatments were analyzed using analysis of variance (ANOVA) and Duncan's Multiple Range Test (DMRT) at 5 % significance level.

RESULTS AND DISCUSSION

Alterations in microbial mineralization and nutrient cycling may control the long-term response of ecosystem to elevated CO₂, because microorganisms are regulators of decomposition, and understanding of microbial activity is so crucial. Elevated CO₂ concentration can affect extracellular enzyme activities in several ways.

In *Rabi* 2013, of the three enzymes examined, dehydrogenase, FDA and urease activities were significantly higher in soils under elevated CO₂ than in ambient. In this study, the increased air temperature in vertisols condition coupled with higher amount of C substrates through rhizodeposits geared up the rate of enzyme activity under elevated CO₂ + temperature condition. In the present study, higher activity of FDA hydrolysis rate noticed in wild oat with 26.5 (µg fluorescein/g/h) at elevated CO₂ - ambient temperature levels. Fluorescein diacetate (FDA) hydrolysis rate are the widely accepted as an accurate method for measuring total microbial activity in aerobic soils. Because they mediated simultaneously by protease, esterase and lipase and thereby reflect the activities of these enzymes in soil (Adam and Duncan 2001). Since 90% of energy flow in a system passes through microbial decomposers and heterotrophic microorganisms are predominant in soil, FDA hydrolysis is thought to reflect overall soil microbiological activity (Schnurer and Rosswall 1982). The activity of dehydrogenase is considered as an indicator of the oxidative metabolism in soils and the microbiological activity, because it is extremely intracellular and able to function within the viable cells.

Dehydrogenase activity was higher at enrichment of CO₂ for *P. minor* with 35.9 (µg TPF/g soil/24 h) followed by wild oat (34.8µg TPF/g soil/24 h). Maximum dehydrogenase activities were found at

the panicle initiation stage and it declined sharply at maturity (Sarathambal *et al.* 2015). Urease the key enzyme hydrolyses urea to ammonia and CO₂ had significantly higher in enrichment of CO₂-elevated temperature with wild oat 34.6 (µg NH₄/g soil/24 h) followed by wheat (31.5 µg NH₄/g soil/24 h) with elevated CO₂ and ambient temperature. The enzyme activities in soil were strongly influenced by exposure to elevated CO₂ as well as by the interactive effect of the elevated CO₂ and temperature throughout the plant growth stages. Moorhead and Linkins (1997) suggested that elevated CO₂ altered the soil enzyme characteristics in a tussock tundra ecosystem.

The microorganisms in soils play a key role in the responses of ecosystem to global climate changes, as they regulate the dynamics of organic matter deposition and plant nutrient availability. However, the increasing atmospheric CO₂ concentrations will indirectly influence microbial populations in soils, through increased root biomass, total rhizodeposition, chemical composition of plant tissues and root exudates probably change when CO₂ is enriched, because CO₂ concentration in soil is much greater than the atmospheric CO₂ (Yue *et al.* 2007).

Table 1. Soil enzyme activity of wheat and associated weed species under OTCs as influenced by elevated CO₂ and temperature

Level of CO ₂ /temperature treatment	Crop / weed	Soil enzyme activity		
		Fluorescein diacetate hydrolysis rate (µg fluorescein/g/h)	Dehydrogenase (µg TPF/g soil/24 h)	Urease (µg NH ₄ /g soil/24 h)
Ambient CO ₂ + ambient temperature	Wheat	18.9 ^{bcd}	25.6 ^{cd}	25.9 ^{bcd}
	<i>P. minor</i>	16.5 ^{de}	24.3 ^d	26.5 ^{bcd}
	Wild oat	14.6 ^{ef}	27.8 ^{cd}	24.6 ^{cd}
Elevated CO ₂ + ambient temperature	Wheat	21.5 ^b	26.9 ^{cd}	31.5 ^{ab}
	<i>P. minor</i>	20.6 ^{bc}	35.9 ^a	20.9 ^d
	Wild oat	26.5 ^a	34.8 ^{ab}	29.5 ^{abc}
Ambient CO ₂ + elevated temperature	Wheat	15.2 ^{def}	22.3 ^d	25.2 ^{cd}
	<i>P. minor</i>	12.3 ^f	23.6 ^d	22.3 ^d
	Wild oat	11.5 ^f	24.5 ^{dg}	21.5 ^d
Elevated CO ₂ + elevated temperature	Wheat	15.6 ^{def}	26.9 ^{cd}	30.6 ^{abc}
	<i>P. minor</i>	16.8 ^{cde}	28.7 ^{bcd}	21.8 ^d
	Wild oat	14.6 ^{ef}	31.5 ^{abc}	34.6 ^a

Values represent mean of three replications and values followed by the same letter in each column are not significantly different from each other as detected by DMRT (p ≤ 0.05)

Table 2. Soil enzyme activity of rice and associated weeds under OTCs as influenced by elevated CO₂ and temperature

Levels of CO ₂ /temperature treatment	Type of cultivar	Soil enzyme activity		
		Fluorescein diacetate hydrolysis rate (µg fluorescein/g/h)	Dehydrogenase (µg TPF/g soil/24 h)	Urease (µg NH ₄ /g soil/24 h)
Ambient CO ₂ + ambient temperature	MRWR27	9.6 ^{ef}	26.5 ^d	39.1 ^{ab}
	W1R4	8.9 ^{efg}	27.6 ^{bcd}	33.9 ^d
	C9	7.6 ^{fg}	18.9 ^g	36.5 ^c
	EC	8.7 ^{efg}	32.4 ^b	36.2 ^c
Elevated CO ₂ + ambient temperature	MRWR27	12.3 ^d	28.6 ^{bc}	39.8 ^c
	W1R4	15.6 ^c	29.8 ^{bc}	36.9 ^c
	C9	11.2 ^{de}	19.5 ^g	42.9 ^{ab}
	EC	18.5 ^{ab}	39.8 ^a	45.6 ^a
Ambient CO ₂ + elevated temperature	MRWR27	6.5 ^g	25.8 ^d	23.6 ^f
	W1R4	9.6 ^{ef}	20.6 ^g	29.8 ^e
	C9	8.7 ^{efg}	22.3 ^e	32.8 ^d
	EC	6.9 ^g	18.9 ^g	33.6 ^d
Elevated CO ₂ + elevated temperature	MRWR27	12.8 ^d	22.3 ^e	34.6 ^d
	W1R4	18.2 ^{ab}	18.9 ^h	29.5 ^e
	C9	16.4 ^{bc}	16.9 ⁱ	28.9 ^e
	EC	19.7 ^a	34.8 ^b	35.5 ^d

Values represent mean of three replications and values followed by the same letter in each column are not significantly different from each other as detected by DMRT (p ≤ 0.05); MRWR27-weedy rice; W1R4-wild rice; C9-cultivated rice; EC- *Echinochloa crusgalli*

Repeated soil sampling during *Kharif* 2014, revealed an enhancement of enzyme activities under elevated CO₂ levels (Table 2). Whereas among the rice associated weeds tested, *Echinochloa crusgalli* rhizosphere responds positively to elevated CO₂ and temperature. In the present study, higher FDA hydrolysis rate noticed in *E. crusgalli* with 19.8 and 18.5 (µg fluorescein/g/h) in elevated CO₂ - elevated temperature and in elevated CO₂ - ambient temperature levels respectively. Dehydrogenase activity was higher at enrichment of CO₂ for *E. crusgalli* with 39.8 (µg TPF/g soil/24 h) followed by elevated CO₂-elevated temperature 34.8 (µg TPF/g soil/24 h). Urease the key enzyme hydrolyses urea to ammonia and CO₂ had significantly higher in enrichment of CO₂-ambient temperature with *E. crusgalli* 45.6 (µg NH₄/g soil/24 h) followed by cultivated rice (C9) 42.9 (µg NH₄/g soil/24 h). Earlier studies conducted by Sardans *et al.* (2008) in soil warming experiment also observed higher enzyme activities under soil warming depending upon the seasonal variations. Similar to our findings, activities of the dehydrogenase, FDA hydrolysis, and urease increased under elevated temperature with CO₂ enrichment resulted in higher microbial activities and increased rate of C mineralization (King *et al.* 2004, Naidu and Sarathambal 2012). A further source of microbial substrates for enzymatic decomposition could be rhizodeposition derived from turnover of fine roots, root hairs and secretion of soluble root exudates; and turnover of rhizosphere-associated microbial biomass. Rhizodeposition was roughly doubled in elevated compared with the ambient CO₂ treatment over the last 4 years of the experiment (Pendall *et al.* 2004). Since rhizodeposition and newly formed roots enlarged the pool of easily available substrates mainly in the weed species like *E. crusgalli*, wild oat, *P. minor*, increased enzyme production in this soil than the crops.

REFERENCES

Adam G and Duncan H. 2001. Development of a sensitive and rapid method for the measurement of total microbial activity using fluorescein diacetate (FDA) in a range of soils. *Soil Biology and Biochemistry* **33**: 943-951.

- Bremner JM and Mulvaney RL. 1978. Urease activity in soils. pp.149-196. In: *Soil enzymes*. (Burns RG Ed.), Academic press, Newyork, USA.
- Casida LEJ, Klein DA and Santoro T. 1964. Soil dehydrogenase activity. *Soil Science* **98**: 370-376.
- Kempers AJ and Zweers A. 1986. Ammonium determination in soil extracts by salicylate method. *Communications in Soil Science and Plant Analysis* **17**: 715-723.
- King JS, Thomas RB and Strain BR. 1997. Morphology and tissue quality of seedling root systems of *Pinus taeda* and *Pinus ponderosa* as affected by varying CO₂, temperature, and nitrogen. *Plant and Soil* **195**: 107-119.
- Kirkpatrick LA and Feenay BC. 2005. *A Simple Guide to SPSS for Windows, for Version 12.0*. Thomson, Wadsworth.
- Moorhead DL and Linkins AE. 1997. Elevated CO₂ alters belowground exoenzyme activities in tussock tundra. *Plant Soil* **189**: 321-329.
- Naidu VSGR and Sarathambal C. 2012. Effect of elevated and atmosphere CO₂ on soil microorganisms, soil respiration and enzyme activities. *Journal of Soil biology and Ecology* **32**: 82-92
- Pendall E, Bridgham S, Hanson P, Hungate B, Kicklighter D and Johnson D. 2004. Belowground process responses to elevated CO₂ and temperature: a discussion of observations, measurement methods and models. *New Phytologist* **162**: 311-322.
- Sarathambal C, Ilamurugu K, Balachandar D, Chinnadurai C and Yogita Gharde. 2015. Characterization and crop production efficiency of diazotrophic isolates from the rhizosphere of semi-arid tropical grasses of India. *Applied Soil Ecology* **87**: 1-10.
- Sardans J, Penuelas J and Estiarte M. 2008. Changes in soil enzymes related to C and N cycle and in soil C and N content under prolonged warming and drought in a Mediterranean shrubland. *Applied Soil Ecology* **39**: 223-235.
- Schnurer, J and T. Rosswall. 1982. Fluorescein dicetate hydrolysis as a measure of total microbial activity in soil and litter. *Applied Environmental Microbiology* **43**: 1256-1261.
- Stager C. 2012. What Happens after Global Warming? *Nature Education Knowledge* **3**(3): 7.
- XLSTAT. 2010. Addinsoft SARL, Paris. Available at <http://www.xlstat.com>.
- Yue J, Shi Y, Zheng X, Huang G and Zhu J. 2007. The influence of free-air CO₂ enrichment on microorganisms in paddy soil in the rice- growing season. *Applied Soil Ecology* **35**: 154-162.