

Induced polyploidy in black pepper (*Piper nigrum* L.)

R. R. NAIR AND P. N. RAVINDRAN
National Research Centre for Spices,
Manikunnu P.O., Calicut - 673 012
Kerala, India.

ABSTRACT

Induced polyploidy in a hybrid cultivar of black pepper Panniyur-I is reported. Open-pollinated seeds were used for polyploidy induction. One tetraploid of Panniyur-I with 104 chromosomes in somatic cells was recovered from plants raised from seeds treated with 0.05% colchicine. The plant was vigorous and possessed all the polyploid syndromes.

Key words : polyploidy, *Piper nigrum*, colchicine.

Piper nigrum L. (Piperaceae) is the source of black pepper of commerce. Somatic chromosome number in most of the cultivated and many of the wild varieties of *P. nigrum* is reported as $2n=52$ (Mathew 1958; Martin & Gregory 1962; Mathew 1973; Samuel & Ravappa 1981; Jose & Sharma 1984; Kada 1986; Samuel & Morawetz 1989). Natural polyploids with $2n=104$ were reported in some wild varieties (Mathew 1958, 1973; Jose & Sharma 1984) and $2n=78$ in a cultivar (Nair, Sasikumar & Ravindran 1993) of *P. nigrum*. But no literature is available so far on induced polyploidy in black pepper. The present study reports the induction of polyploidy in black pepper by treating seeds with colchicine.

were treated with 0.05% and 0.1% colchicine in distilled water for 4 h from 10 am to 1 pm by soaking the seeds in it. Subsequently, the treated seeds were washed in distilled water and kept in moist filter paper in a petridish. The seeds were sown after 20 h. Fifty untreated seeds formed the control. After germination, the seedlings were allowed to grow in pots for 6 months and then transferred to multiplication nursery. In the nursery some of the treated plants showed variation in growth and morphological characters such as stunting, thick leaves, change in leaf shape and stem thickness. These plants were cytologically analysed along with controls.

For cytological analysis, actively grow-

ripened seeds of hybrid black pepper

form. Staining was done with 2% lactopropionic-orcein. Only intact mitotic metaphase plates with reasonably good spread were counted for chromosome number. To determine stomatal frequency, uniform sized leaf discs of both Panniyur-I and induced polyploid were cut and epidermal layers separated by boiling in conc. HNO_3 . Three leaf discs in each case were counted for number of stomata and number of stomata per unit area was also calculated.

On chromosome analysis, it was found that plant No. 75 which was derived

from a seed treated with 0.05% colchicine had a chromosome number of 104 in its somatic cells (Fig. 1 d). Out of 10 mitotic metaphase plates of plant No. C.5 from 10 slides counted, all had 104 chromosomes. The plant was characterized by thick, typically cordate leaves, thick stem, purple shoot tip, stipules and vigorous growth in nursery. The number of stomata per unit area was 95 in the case of Panniyur-I and 104 in induced polyploid. On visual comparison it was found that the stomata in induced polyploid was larger than in Panniyur-I. The plant was strikingly




Fig. 1. Normal and induced polyploid in black pepper (var. Panniyur-I)
a. A normal plant of Panniyur-I b. Induced polyploid from colchicine treated seed
c. A mitotic metaphase cell of Panniyur-I showing $2n=52$ d. A mitotic metaphase cell of induced polyploid showing $2n=104$ (Bar represents $10 \mu\text{m}$)

different in shoot tip colour from Panniyur-I, which has greenish white shoot tip. Since shoot tip colour of all other plants including controls were greenish white it is possible that the colchipsoid mentioned may have been derived from an out-crossed seed. Production of greenish white shoot tipped progenies alone by Panniyur-I has already been reported by Ibrahim, Pillai & Sasikumar (1985). According to Ravindran, Sasikumar & Babu (1982) pigmentation in black pepper is controlled by two complementary genes A_1 and A_2 and Panniyur-I has a genotype of $A_1a_1a_2a_2$ which will produce only greenish white progenies on selfing.

The present report on induced polyploidy in black pepper is the first of its kind, and gives us hope for production of aneuploids in black pepper. The possibility of an induced polyloid to emerge itself as a high yielding variety cannot also be ruled out.

References

- Ibrahim K K, Sukumara Pillai V & Sasikumar S 1985 Inheritance of anthocyanin pigmentation on stipules in black pepper. *Indian Cocoa, Arecanut and Spices J.* 9 (1) : 12-14.
- Muse J & Sharma A K 1984 Chromosome studies in genus *Piper* L. *J Indian Bot. Soc.* 63:313-319.

Martin F W & Gregory L E 1962 Mode of pollination and factors affecting fruit set in *Piper nigrum* L. in Puerto Rico. *Crop Sci.* 2:295-299.

Mathew P M 1958 Studies on Piperaceae. *J. Indian Bot. Soc.* 37:165-171.

Mathew P M 1971 Karyomorphological studies in *Piper nigrum* L. *J. Plant. Crops* 1 (Suppl) :15-18.

Nair R R, Sasikumar B & Ravindran P N 1993 Polyploidy in a cultivar of black pepper and its open pollinated progenies. *Cytologia* (in press).

Okada H 1986 Karyomorphology and relationships in some genera of Sauraceae and Piperaceae. *Bot. Mag. Tokyo* 99:289-299.

Ravindran P N, Sasikumar B & Nirmal Babu K 1992 Genetics of shoot-tip colour in black pepper. *J. Plant. Crops* 20:76-78.

Samuel M R A & Bavappa K V A 1981 Chromosome number in the genus *Piper*. *Curr. Sci.* 50:197-198.

Samuel R & Morawetz W 1989 Chromosomal evolution within Piperaceae. *Plant. Syst. Evoln.* 166:105-117.